arXiv:1606.08327v2 [math.NT] 11 Jul 2016

preprint: July 11, 2016

A kind of orthogonal polynomials and related identities

Zhi-Hong Sun

School of Mathematical Sciences
Huaiyin Normal University
Huaian, Jiangsu 223001, P.R. China
zhsun@hytc.edu.cn
http://www.hytc.edu.cn/xsjl/szh

Abstract In this paper we investigate the properties of a kind of orthogonal polynomials and related identities. As applications we extend recent work of V.J.W. Guo.

Keywords: polynomial; identity; three-term recurrence

Mathematics Subject Classification 2010: Primary 11B83, Secondary 05A10, 05A19, 11A07, 30B10, 33B15, 33C47

1. Introduction

Let n be a nonnegative integer. By [G, (3.17)],

$$\sum_{k=0}^n \binom{n}{k}\binom{x}{k}t^k = \sum_{k=0}^n \binom{n}{k}\binom{x+k}{n}(t-1)^{n-k}. \tag{1.1}$$

Define

$$d_n(x) = \sum_{k=0}^n \binom{n}{k}\binom{x}{k}2^k.$$

For given positive integers m and n, $d_n(m)$ is the number of lattice paths from $(0,0)$ to (m,n), with jumps $(0,1)$, $(1,1)$ or $(1,0)$. $\{d_n(m)\}$ are called Delannoy numbers. See [BS]. In [Su] Z.W. Sun deduced some supercongruences involving $d_n(x)$. Actually, he obtained congruences for

$$\sum_{k=0}^{p-1} d_k(x)^2,\ \sum_{k=0}^{p-1}(-1)^k d_k(x)^2,\ \sum_{k=0}^{p-1}(2k+1)d_k(x)^2 \quad\text{and}\quad \sum_{k=0}^{p-1}(-1)^k(2k+1)d_k(x)^2$$

modulo p^2, where p is a odd prime and x is a p-adic integer. Let $\mathbb{Z}$ be the set of integers and let $\mathbb{Z}^+$ be the set of positive integers. Z.W. Sun also conjectured that for any $n\in\mathbb{Z}^+$ and $x\in\mathbb{Z}$,

$$x(x+1)\sum_{k=0}^{n-1}(2k+1)d_k(x)^2 \equiv 0 \pmod{2n^2} \tag{1.2}$$

and

$$\sum_{k=0}^{n-1}\varepsilon^k(2k+1)d_k(x)^{2m}\equiv 0\ (\mathrm{mod}\ n)\quad \text{for given } \varepsilon\in\{1,-1\} \text{ and } m\in\mathbb{Z}^+. \tag{1.3}$$

Recently, Guo[Gu] proved the above two congruences by using the identity

$$d_n(x)^2=\sum_{k=0}^{n}\binom{n+k}{2k}\binom{x}{k}\binom{x+k}{k}4^k. \tag{1.4}$$

Guo proved (1.4) by using Maple and Zeilberger's algorithm, and W. Zudilin stated that (1.4) can be deduced from two transformation formulas for hypergeometric series. See [Gu] and [Sl, (1.7.1.3) and (2.5.32)].

Let $\{p_n(x)\}$ be the polynomials satisfying

$$p_{-1}(x)=0,\ p_0(x)=1\quad\text{and}\quad xp_n(x)=A_np_{n+1}(x)+B_np_n(x)+C_np_{n-1}(x)(n\ge 0). \tag{1.5}$$

From [GR, pp.199-200] or [EMOT, p.159] we have the following Christoffel-Darboux formulas:

$$(x-y)\sum_{k=0}^{n}v_kp_k(x)p_k(y)=A_nv_n(p_{n+1}(x)p_n(y)-p_n(x)p_{n+1}(y)), \tag{1.6}$$

$$\sum_{k=0}^{n}v_kp_k(x)^2=A_nv_n(p'_{n+1}(x)p_n(x)-p'_n(x)p_{n+1}(x)), \tag{1.7}$$

where $v_n=\frac{A_0A_1\cdots A_n}{C_1\cdots C_n}$ and $f'(x)$ is the derivative of $f(x)$. In Section 2 we extend Christoffel-Darboux formulas for three-term recurrence sequences.

In Section 3 we introduce the polynomials

$$d_n^{(r)}(x)=\sum_{k=0}^{n}\binom{x+r+k}{k}\binom{x-r}{n-k}\ (n=0,1,2,\ldots) \tag{1.8}$$

and investigate the properties of $d_n^{(r)}(x)$. Clearly $d_n(x)=d_n^{(0)}(x)$ by (1.1). Thus, $d_n^{(r)}(x)$ is a generalization of $d_n(x)$. We establish many identities for $d_n^{(r)}(x)$. As examples, we have

$$d_n^{(r)}(x)=d_n^{(r+1)}(x)-d_{n-2}^{(r+1)}(x), \tag{1.9}$$

$$(2r+1)\sum_{k=0}^{n-1}(2k+2r+1)d_k^{(r)}(x)^2=n^2d_n^{(r)}(x)^2-4(x-r)(x+1+r)d_{n-1}^{(r+1)}(x)^2, \tag{1.10}$$

and for $r\in\{0,1,2,\ldots\}$,

$$\sum_{k=0}^{n-1}(-1)^k(2k+2r+1)\frac{k!d_k^{(r)}(x)^2}{(k+2r)!}=\frac{(-1)^nn!}{(n+2r-1)!}\big(nd_n^{(r)}(x)^2-(n+1)d_{n-1}^{(r)}(x)d_{n+1}^{(r)}(x)\big). \tag{1.11}$$

Using (1.10) we prove that for $r \in \{0,1,2,\ldots\}$, $n \in \mathbb{Z}^+$ and $x \in \mathbb{Z}$,

$$(2r+1)\prod_{k=-r}^{r}(x+k)(x+1-k)\cdot\sum_{k=0}^{n-1}(2k+2r+1)d_k^{(r)}(x)^2 \equiv 0 \pmod{2n^2(n+1)^2\cdots(n+2r)^2}.$$

Taking $r=0$ in the congruence yields (1.2). Taking $r=0$ in (1.11) yields (1.3) in the case $\varepsilon=-1$ and $m=1$. In Section 3 we also extend (1.4) by showing that

$$d_n^{(r)}(x)^2 = \binom{n+2r}{n}\sum_{m=0}^{n}\frac{\binom{x-r}{m}\binom{x+r+m}{m}\binom{n+2r+m}{n-m}}{\binom{m+2r}{m}}4^m \quad \text{for} \quad r \notin \{-\frac{1}{2},-\frac{2}{2},-\frac{3}{2},\ldots\}. \tag{1.12}$$

We also state that the linearization of $d_m^{(r)}(x)d_n^{(r)}(x)$:

$$d_m^{(r)}(x)d_n^{(r)}(x) = \sum_{k=0}^{\min\{m,n\}}\binom{m+n-2k}{m-k}\binom{2r+m+n-k}{k}(-1)^k d_{m+n-2k}^{(r)}(x). \tag{1.13}$$

In Section 4 we introduce the orthogonal polynomials $\{D_n^{(r)}(x)\}$, which are related to $\{d_n^{(r)}(x)\}$. Let $\{D_n^{(r)}(x)\}$ be the polynomials given by

$$D_0^{(r)}(x)=1,\ D_1^{(r)}(x)=x \quad \text{and} \quad D_{n+1}^{(r)}(x)=xD_n^{(r)}(x)-n(n+2r)D_{n-1}^{(r)}(x)\ (n\ge 1).$$

We state that $D_n^{(r)}(x)=(-i)^n n!d_n^{(r)}(\frac{ix-1}{2})$. By [GR, p.175], $\{D_n^{(r)}(x)\}$ are orthogonal polynomials for $r>-\frac{1}{2}$, although we have not found their weight functions. We obtain some properties of $\{D_n^{(r)}(x)\}$. In particular, we show that

$$D_n^{(r)}(x)^2 - D_{n+1}^{(r)}(x)D_{n-1}^{(r)}(x) > 0 \quad \text{for } r > -\frac{1}{2} \text{ and real } x. \tag{1.14}$$

In Section 5 we discuss the orthogonal polynomials

$$C_n(x)=\sum_{k=0}^{n}\binom{n}{k}\frac{1}{k!^2}\prod_{j=0}^{k-1}(x-j(j+1)).$$

2. Simple identities for three-term recurrence sequences

For given sequences $\{a_n\}$, $\{b_n\}$ and constant c with $a_n c \neq 0$ $(n=1,2,\ldots)$ let $\{s_n\}$ be the sequence defined by

$$s_0=1,\ s_1=\frac{b_0}{a_1} \quad \text{and} \quad a_{n+1}s_{n+1}=b_n s_n + c a_n s_{n-1}\ (n\ge 1). \tag{2.1}$$

Theorem 2.1. *Let $\{s_n\}$ be the sequence satisfying (2.1). Then*

$$\sum_{m=0}^{n-1} b_m c^{n-1-m} s_m^2 = a_n s_n s_{n-1} \quad \textit{for} \quad n=1,2,3,\ldots.$$

Proof. From the definition of $\{s_n\}$ we see that

$$\frac{b_m}{c^m}s_m^2 = \frac{a_{m+1}}{c^m}s_{m+1}s_m - \frac{a_m}{c^{m-1}}s_m s_{m-1}.$$

Thus,

$$\sum_{m=1}^{n-1}\frac{b_m}{c^m}s_m^2 = \sum_{m=1}^{n-1}\left(\frac{a_{m+1}}{c^m}s_{m+1}s_m - \frac{a_m}{c^{m-1}}s_m s_{m-1}\right) = \frac{a_n}{c^{n-1}}s_n s_{n-1} - a_1 s_1 s_0.$$

This yields the result.

Corollary 2.1. *Let $\{P_n(x)\}$ be the Legendre polynomials given by*

$$P_0(x) = 1,\ P_1(x) = x \quad \textit{and} \quad (n+1)P_{n+1}(x) = (2n+1)xP_n(x) - nP_{n-1}(x) \quad (n \geq 1).$$

Then

$$\sum_{m=0}^{n-1}(-1)^{n-1-m}(2m+1)P_m(x)^2 = n\frac{P_n(x)P_{n-1}(x)}{x} \quad \textit{for } n \in \mathbb{Z}^+ \textit{ and } x \neq 0.$$

Proof. Taking $a_n = n$, $b_n = (2n+1)x$ and $c = -1$ in Theorem 2.1 yields the result.

Corollary 2.2. *Let $\{A_n\}$ be the Apéry numbers given by*

$$A_n = \sum_{k=0}^{n}\binom{n}{k}^2\binom{n+k}{k}^2 \ (n \geq 0).$$

Then

$$\sum_{m=0}^{n-1}(-1)^m(2m+1)(17m^2+17m+5)A_m^2 = (-1)^{n-1}n^3A_nA_{n-1}.$$

Proof. In 1987 Beukers[Be] showed that

$$(n+1)^3A_{n+1} = (2n+1)(17n^2+17n+5)A_n - n^3A_{n-1} \quad (n \geq 1).$$

This can be easily checked by Maple. Now taking $a_n = n^3$, $b_n = (2n+1)(17n^2+17n+5)$ and $c = -1$ in Theorem 2.1 yields the result.

Corollary 2.3. *Let $\{A'_n\}$ be given by*

$$A'_n = \sum_{k=0}^{n}\binom{n}{k}^2\binom{n+k}{k} \ (n \geq 0).$$

Then

$$\sum_{m=0}^{n-1}(11m^2+11m+3){A'}_m^2 = n^2A'_nA'_{n-1}.$$

Proof. It is well known that

$$(n+1)^2A'_{n+1} = (11n^2+11n+3)A'_n + n^2A'_{n-1} \ (n \geq 1).$$

Now taking $a_n = n^2$, $b_n = 11n^2 + 11n + 3$ and $c = 1$ in Theorem 2.1 yields the result.

Corollary 2.4. *Let $\{f_n\}$ be the Franel numbers given by*

$$f_n = \sum_{k=0}^{n} \binom{n}{k}^3 \ (n \geq 0).$$

Then

$$\sum_{m=0}^{n-1} (7m^2 + 7m + 2) 8^{n-1-m} f_m^2 = n^2 f_n f_{n-1}.$$

Proof. It is well known that

$$(n+1)^2 f_{n+1} = (7n^2 + 7n + 2) f_n + 8n^2 f_{n-1} \ (n \geq 1).$$

Now taking $a_n = n^2$, $b_n = 7n^2 + 7n + 2$ and $c = 8$ in Theorem 2.1 yields the result.

Corollary 2.5. *Let $\{S_n\}$ be given by*

$$S_n = \sum_{k=0}^{n} \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \ (n \geq 0).$$

Then

$$\sum_{m=0}^{n-1} (3m^2 + 3m + 1)(-32)^{n-1-m} S_m^2 = \frac{n^2 S_n S_{n-1}}{4}.$$

Proof. Using Maple we find that

$$(n+1)^2 S_{n+1} = 4(3n^2 + 3n + 1) S_n - 32n^2 S_{n-1} \ (n \geq 1).$$

Now taking $a_n = n^2$, $b_n = 4(3n^2 + 3n + 1)$ and $c = -32$ in Theorem 2.1 yields the result.

Corollary 2.6. *Let $\{d_n(x,t)\}$ be given by*

$$d_n(x,t) = \sum_{k=0}^{n} \binom{n}{k} \binom{x}{k} t^k \ (n \geq 0).$$

Then

$$\sum_{m=0}^{n-1} (1 + tx + (2-t)m)(t-1)^{n-1-m} d_m(x,t)^2 = n d_n(x,t) d_{n-1}(x,t).$$

Proof. Using Maple we find that

$$(n+1) d_{n+1}(x,t) = (1 + tx + (2-t)n) d_n(x,t) + (t-1) n d_{n-1}(x,t) \ (n \geq 1)$$

Now taking $a_n = n$, $b_n = 1 + tx + (2-t)n$ and $c = t - 1$ in Theorem 2.1 yields the result.

Remark 2.1 For $\{A_n\}, \{A'_n\}, \{S_n\}$ and $\{f_n\}$ see A005259, A005258, A053175 and A000172 in Sloane's database "The On-Line Encyclopedia of Integer Sequences".

For given sequences $\{a_n\}$, $\{b_n\}$, $\{b'_n\}$ and constant c with $b_0 = a_1$, $b'_0 = 0$ and $a_n c \neq 0$ ($n = 1, 2, \ldots$) let $\{s_n(x)\}$ be the sequence satisfying

$$s_0(x) = 1 \quad \text{and} \quad a_{n+1}s_{n+1}(x) = (b_n s_1(x) + b'_n)s_n(x) + ca_n s_{n-1}(x) \ (n \geq 1). \tag{2.2}$$

Then

$$a_{m+1}s_{m+1}(x)s_m(y) = (b_m s_1(x) + b'_m)s_m(x)s_m(y) + ca_m s_{m-1}(x)s_m(y).$$

Set $s_n(x) = (\sqrt{c})^n s'_n(x)$. Then

$$(\sqrt{c})^{2m+1}a_{m+1}s'_{m+1}(x)s'_m(y) = (b_m s_1(x) + b'_m)s_m(x)s_m(y) + (\sqrt{c})^{2m-1}ca_m s'_{m-1}(x)s'_m(y).$$

Thus,

$$\begin{aligned}
&(\sqrt{c})^{2m+1}a_{m+1}(s'_{m+1}(x)s'_m(y) \pm s'_m(x)s'_{m+1}(y)) \\
&= (b_m s_1(x) + b'_m \pm (b_m s_1(y) + b'_m))s_m(x)s_m(y) + (\sqrt{c})^{2m+1}a_m(s'_{m-1}(x)s'_m(y) \pm s'_m(x)s'_{m-1}(y)).
\end{aligned}$$

That is,

$$\begin{aligned}
&\frac{(s_1(x) \pm s_1(y))b_m + b'_m \pm b'_m}{(\sqrt{c})^{2m+1}}s_m(x)s_m(y) \\
&= a_{m+1}(s'_{m+1}(x)s'_m(y) \pm s'_m(x)s'_{m+1}(y)) - a_m(s'_{m-1}(x)s'_m(y) \pm s'_m(x)s'_{m-1}(y)) \\
&= a_{m+1}(s'_{m+1}(x)s'_m(y) \pm s'_m(x)s'_{m+1}(y)) \mp a_m(s'_m(x)s'_{m-1}(y) \pm s'_{m-1}(x)s'_m(y)).
\end{aligned}$$

Hence

$$\begin{aligned}
&\sum_{m=1}^{n-1} \frac{(s_1(x) + s_1(y))b_m + 2b'_m}{(\sqrt{c})^{2m+1}}s_m(x)s_m(y) \\
&= \sum_{m=1}^{n-1} \big(a_{m+1}(s'_{m+1}(x)s'_m(y) + s'_m(x)s'_{m+1}(y)) - a_m(s'_m(x)s'_{m-1}(y) + s'_{m-1}(x)s'_m(y)\big) \\
&= a_n(s'_n(x)s'_{n-1}(y) + s'_{n-1}(x)s'_n(y)) - a_1(s'_1(x)s'_0(y) + s'_0(x)s'_1(y)) \\
&= \frac{a_n(s_n(x)s_{n-1}(y) + s_{n-1}(x)s_n(y))}{(\sqrt{c})^{2n-1}} - a_1\frac{s_1(x) + s_1(y)}{\sqrt{c}}
\end{aligned}$$

and

$$\begin{aligned}
&\sum_{m=1}^{n-1}(-1)^m\frac{(s_1(x) - s_1(y))b_m}{(\sqrt{c})^{2m+1}}s_m(x)s_m(y) \\
&= \sum_{m=1}^{n-1}(-1)^m\big(a_{m+1}(s'_{m+1}(x)s'_m(y) - s'_m(x)s'_{m+1}(y)) + a_m(s'_m(x)s'_{m-1}(y) - s'_{m-1}(x)s'_m(y)\big) \\
&= (-1)^{n-1}a_n(s'_n(x)s'_{n-1}(y) - s'_{n-1}(x)s'_n(y)) - a_1(s'_1(x)s'_0(y) - s'_0(x)s'_1(y)) \\
&= (-1)^{n-1}\frac{a_n(s_n(x)s_{n-1}(y) - s_{n-1}(x)s_n(y))}{(\sqrt{c})^{2n-1}} - a_1\frac{s_1(x) - s_1(y)}{\sqrt{c}}.
\end{aligned}$$

This yields the following theorem.

Theorem 2.2. *Let $\{s_n(x)\}$ be given by (2.2). For $n \in \mathbb{Z}^+$ we have*

$$\sum_{m=0}^{n-1}((s_1(x)+s_1(y))b_m+2b'_m)c^{n-1-m}s_m(x)s_m(y)=a_n(s_n(x)s_{n-1}(y)+s_{n-1}(x)s_n(y))$$

and

$$\sum_{m=0}^{n-1}b_m(-c)^{n-1-m}s_m(x)s_m(y)=\frac{a_n(s_n(x)s_{n-1}(y)-s_{n-1}(x)s_n(y))}{s_1(x)-s_1(y)}.$$

Theorem 2.3. *Let $\{d_n(x,t)\}$ be given by*

$$d_n(x,t)=\sum_{k=0}^{n}\binom{n}{k}\binom{x}{k}t^k\ (n\geq 0).$$

Then for any positive integer n,

$$\sum_{m=0}^{n-1}(1-t)^{n-1-m}d_m(x,t)d_m(y,t)=\frac{n(d_n(x,t)d_{n-1}(y,t)-d_{n-1}(x,t)d_n(y,t))}{t(x-y)}$$

and

$$\begin{aligned}&\sum_{m=0}^{n-1}(2+t(x+y)+(4-2t)m)(t-1)^{n-1-m}d_m(x,t)d_m(y,t)\\&=n(d_n(x,t)d_{n-1}(y,t)+d_{n-1}(x,t)d_n(y,t)).\end{aligned}$$

Proof. As $d_1(x,t)=1+tx$ and

$$(n+1)d_{n+1}(x,t)=(1+tx+(2-t)n)d_n(x,t)+(t-1)nd_{n-1}(x,t)\ (n\geq 1),$$

putting $a_n=n$, $b_n=1$, $b'_n=(2-t)n$, $c=t-1$ and $s_n(x)=d_n(x,t)$ in Theorem 2.2 gives the result.

3. The polynomials $\{d_n^{(r)}(x)\}$

Recall that

$$d_n(x)=\sum_{k=0}^{n}\binom{n}{k}\binom{x}{k}2^k\ (n=0,1,2,\ldots).$$

By (1.1),

$$d_n(x)=\sum_{k=0}^{n}\binom{n}{k}\binom{x+k}{n}=\sum_{k=0}^{n}\binom{x+k}{k}\binom{x}{n-k}. \tag{3.1}$$

The first few $\{d_n(x)\}$ are shown below:

$$d_0(x)=1,\ d_1(x)=2x+1,\ d_2(x)=2x^2+2x+1,\ d_3(x)=\frac{4}{3}x^3+2x^2+\frac{8}{3}x+1,$$
$$d_4(x)=\frac{2}{3}x^4+\frac{4}{3}x^3+\frac{10}{3}x^2+\frac{8}{3}x+1,\ d_5(x)=\frac{4}{15}x^5+\frac{2}{3}x^4+\frac{8}{3}x^3+\frac{10}{3}x^2+\frac{46}{15}x+1.$$

Taking $t=2$ in Corollary 2.6 we get

$$\sum_{m=0}^{n-1} d_m(x)^2=\frac{nd_n(x)d_{n-1}(x)}{1+2x}. \tag{3.2}$$

Taking $t=2$ in Theorem 2.3 we obtain

$$\sum_{m=0}^{n-1} d_m(x)d_m(y)=\frac{n(d_n(x)d_{n-1}(y)+d_{n-1}(x)d_n(y))}{2(1+x+y)} \tag{3.3}$$

and

$$\sum_{m=0}^{n-1}(-1)^{n-1-m}d_m(x)d_m(y)=\frac{n(d_n(x)d_{n-1}(y)-d_{n-1}(x)d_n(y))}{2(x-y)}. \tag{3.4}$$

Now we introduce the following generalization of $\{d_n(x)\}$.

Definition 3.1. *Let $\{d_n^{(r)}(x)\}$ be the polynomials given by*

$$d_n^{(r)}(x)=\sum_{k=0}^{n}\binom{x+r+k}{k}\binom{x-r}{n-k}\ (n=0,1,2,\ldots).$$

For convenience we also define $d_{-1}^{(r)}(x)=0$.

As $\binom{-a}{k}=(-1)^k\binom{a+k-1}{k}$ we see that

$$d_n^{(r)}(x)=\sum_{k=0}^{n}\binom{-1-x-r}{k}(-1)^k\binom{x-r}{n-k}=\sum_{k=0}^{n}\binom{-1-x-r}{n-k}(-1)^{n-k}\binom{x-r}{k}.$$

Hence

$$d_n^{(r)}(-1-x)=(-1)^n d_n^{(r)}(x). \tag{3.5}$$

It is easily seen that for any nonnegative integers k,n and r with $k\le n$,

$$\binom{x+r}{2r}\binom{x+r+k}{k}\binom{x-r}{n-k}=\binom{n+2r}{2r}\binom{n}{k}\binom{x+r+k}{n+2r}.$$

Thus,

$$\binom{x+r}{2r}d_n^{(r)}(x)=\binom{n+2r}{2r}\sum_{k=0}^{n}\binom{n}{k}\binom{x+r+k}{n+2r}\quad\text{for}\quad r=0,1,2,\ldots. \tag{3.6}$$

By (3.1), $d_n(x) = d_n^{(0)}(x)$. The first few $\{d_n^{(r)}(x)\}$ are shown below:

$$d_0^{(r)}(x) = 1,\ d_1^{(r)}(x) = 2x+1,\ d_2^{(r)}(x) = 2x^2+2x+r+1,$$
$$d_3^{(r)}(x) = \frac{4}{3}x^3 + 2x^2 + \big(2r+\frac{8}{3}\big)x + r + 1,$$
$$d_4^{(r)}(x) = \frac{2}{3}x^4 + \frac{4}{3}x^3 + \big(2r+\frac{10}{3}\big)x^2 + \big(2r+\frac{8}{3}\big)x + \frac{(r+1)(r+2)}{2},$$
$$d_5^{(r)}(x) = \frac{4}{15}x^5 + \frac{2}{3}x^4 + \big(\frac{4}{3}r+\frac{8}{3}\big)x^3 + \big(2r+\frac{10}{3}\big)x^2 + \big(r^2+\frac{11}{3}r+\frac{46}{15}\big)x + \frac{(r+1)(r+2)}{2}.$$

Theorem 3.1. *For $|t|<1$ we have*

$$\sum_{n=0}^{\infty} d_n^{(r)}(x)t^n = \frac{(1+t)^{x-r}}{(1-t)^{x+r+1}}.$$

Proof. By Newton's binomial theorem,

$$\begin{aligned}
&(1+t)^{x-r}(1-t)^{-x-r-1}\\
&= \Big(\sum_{m=0}^{\infty}\binom{x-r}{m}t^m\Big)\Big(\sum_{k=0}^{\infty}\binom{-x-r-1}{k}(-1)^k t^k\Big)\\
&= \sum_{n=0}^{\infty}\Big(\sum_{k=0}^{n}\binom{-x-r-1}{k}(-1)^k\binom{x-r}{n-k}\Big)t^n = \sum_{n=0}^{\infty}\Big(\sum_{k=0}^{n}\binom{x+r+k}{k}\binom{x-r}{n-k}\Big)t^n\\
&= \sum_{n=0}^{\infty} d_n^{(r)}(x)t^n.
\end{aligned}$$

This proves the theorem.

Corollary 3.1. *For any positive integer n we have*

$$d_n^{(r)}\Big(-\frac{1}{2}\Big) = \begin{cases} 0 & \text{if } 2\nmid n,\\ \binom{-\frac{1}{2}-r}{\frac{n}{2}}(-1)^{\frac{n}{2}} & \text{if } 2\mid n.\end{cases}$$

Proof. By Theorem 3.1 and Newton's binomial theorem, for $|t|<1$ we have

$$\sum_{n=0}^{\infty} d_n^{(r)}\big(-\frac{1}{2}\big)t^n = (1-t^2)^{-\frac{1}{2}-r} = \sum_{k=0}^{\infty}\binom{-\frac{1}{2}-r}{k}(-1)^k t^{2k}.$$

Now comparing the coefficients of t^n on both sides yields the result.

Theorem 3.2. *For any positive integer n we have*

$$(n+1)d_{n+1}^{(r)}(x) = (1+2x)d_n^{(r)}(x) + (n+2r)d_{n-1}^{(r)}(x).$$

Proof. By Theorem 3.1, for $|t|<1$ it is clear that

$$\sum_{n=0}^{\infty}(n+1)d_{n+1}^{(r)}(x)t^n - \sum_{n=0}^{\infty} n d_{n-1}^{(r)}(x)t^n$$

$$
\begin{aligned}
&= \Big(\sum_{n=0}^{\infty} d_{n+1}^{(r)}(x)t^{n+1}\Big)' - t\Big(\sum_{n=1}^{\infty} d_{n-1}^{(r)}(x)t^{n}\Big)' \\
&= \big((1+t)^{x-r}(1-t)^{-x-r-1}\big)' - t\big(t(1+t)^{x-r}(1-t)^{-x-r-1}\big)' \\
&= \big((1+t)^{x-r}(1-t)^{-x-r-1}\big)' - t((1+t)^{x-r}(1-t)^{-x-r-1} + t((1+t)^{x-r}(1-t)^{-x-r-1})') \\
&= (1-t^2)\big((x-r)(1+t)^{x-r-1}(1-t)^{-x-r-1} + (1+t)^{x-r}(x+r+1)(1-t)^{-x-r-2}\big) \\
&\quad - t(1+t)^{x-r}(1-t)^{-x-r-1} \\
&= (1+2x+2rt)(1+t)^{x-r}(1-t)^{-x-r-1} \\
&= (1+2x)\sum_{n=0}^{\infty} d_n^{(r)}(x)t^n + 2r\sum_{n=1}^{\infty} d_{n-1}^{(r)}(x)t^n.
\end{aligned}
$$

Now comparing the coefficients of t^n on both sides gives the result.

Remark 3.1 By Theorem 3.2,

$$
\frac{d_{n+1}^{(1)}(x)}{n+2} = \frac{2x+1}{(n+1)(n+2)}d_n^{(1)}(x) + \frac{d_{n-1}^{(1)}(x)}{n+1} \ (n \geq 1),
$$

taking $a_n = \frac{1}{n+1}$, $b_n = \frac{1}{(n+1)(n+2)}$, $b_n' = 0$ and $c = 1$ in Theorem 2.2 gives

$$
2(1+x+y)\sum_{m=0}^{n-1} \frac{d_m^{(1)}(x)d_m^{(1)}(y)}{(m+1)(m+2)} = \frac{d_n^{(1)}(x)d_{n-1}^{(1)}(y) + d_{n-1}^{(1)}(x)d_n^{(1)}(y)}{n+1} \tag{3.7}
$$

and

$$
\sum_{m=0}^{n-1}(-1)^{n-1-m}\frac{d_m^{(1)}(x)d_m^{(1)}(y)}{(m+1)(m+2)} = \frac{d_n^{(1)}(x)d_{n-1}^{(1)}(y) - d_{n-1}^{(1)}(x)d_n^{(1)}(y)}{2(n+1)(x-y)}. \tag{3.8}
$$

Theorem 3.3. *Let n be a nonnegative integer. Then*

$$
d_n^{(r)}(x) = \sum_{k=0}^{[n/2]} \binom{r-1+k}{k} d_{n-2k}(x) = \sum_{k=0}^{n} \binom{2r-1+k}{k} d_{n-k}(x-r)
$$

and

$$
d_n(x) = \sum_{k=0}^{[n/2]} \binom{r}{k}(-1)^k d_{n-2k}^{(r)}(x) = \sum_{k=0}^{n} \binom{2r}{k}(-1)^k d_{n-k}^{(r)}(x+r),
$$

where $[a]$ is the greatest integer not exceeding a.

Proof. As

$$
\sum_{n=0}^{\infty} d_n^{(r)}(x)t^n = (1-t^2)^{-r}\cdot\frac{1}{1-t}\Big(\frac{1+t}{1-t}\Big)^x = (1-t)^{-2r}\cdot\frac{1}{1-t}\Big(\frac{1+t}{1-t}\Big)^{x-r},
$$

we see that

$$
\sum_{n=0}^{\infty} d_n^{(r)}(x)t^n = (1-t^2)^{-r}\sum_{n=0}^{\infty} d_n(x)t^n = (1-t)^{-2r}\sum_{n=0}^{\infty} d_n(x-r)t^n
$$

and so

$$\sum_{n=0}^{\infty} d_n(x)t^n = (1-t^2)^r \sum_{n=0}^{\infty} d_n^{(r)}(x)t^n = (1-t)^{2r} \sum_{n=0}^{\infty} d_n^{(r)}(x+r)t^n.$$

Since $\binom{-a}{k} = (-1)^k\binom{a+k-1}{k}$, applying Newton's binomial theorem and comparing the coefficients of t^n on both sides we deduce the result.

Corollary 3.2. *Let n be a nonnegative integer. Then*

$$d_n^{(r)}(0) = \binom{r+\left[\frac{n}{2}\right]}{\left[\frac{n}{2}\right]}.$$

Proof. Since $\binom{-x}{k} = (-1)^k\binom{x+k-1}{k}$ and $d_n(0) = \sum_{k=0}^{n}\binom{n}{k}\binom{0}{k}2^k = 1$, applying Theorem 3.3 we get

$$\begin{aligned} d_n^{(r)}(0) &= \sum_{k=0}^{[n/2]} \binom{r-1+k}{k} = \sum_{k=0}^{[n/2]} \binom{-r}{k}(-1)^k = \sum_{k=0}^{[n/2]} \left((-1)^k\binom{-r-1}{k} - (-1)^{k-1}\binom{-r-1}{k-1}\right) \\ &= (-1)^{\left[\frac{n}{2}\right]}\binom{-r-1}{\left[\frac{n}{2}\right]} = \binom{r+\left[\frac{n}{2}\right]}{\left[\frac{n}{2}\right]}. \end{aligned}$$

This proves the corollary.

Lemma 3.1. *Let $n \in \mathbb{Z}^+$ and $k \in \{1, 2, \ldots, n\}$. Then*

$$[x^{n-k}]d_n^{(r)}(x) = \sum_{m=k}^{n} \frac{2^{n-m}\cdot(m-1)!}{n!}\Big([x^{m-k}]d_{m-1}^{(r)}(x) + (m-1+2r)[x^{m-k}]d_{m-2}^{(r)}(x)\Big),$$

where $[x^s]f(x)$ is the coefficient of x^s in the power series expansion of $f(x)$.

Proof. By Theorem 3.2, $md_m^{(r)}(x) = (1+2x)d_{m-1}^{(r)}(x) + (m-1+2r)d_{m-2}^{(r)}(x)$. Thus,

$$m[x^{m-k}]d_m^{(r)}(x) = [x^{m-k}]d_{m-1}^{(r)}(x) + 2[x^{m-k-1}]d_{m-1}^{(r)}(x) + (m-1+2r)[x^{m-k}]d_{m-2}^{(r)}(x).$$

It then follows that

$$\begin{aligned} &[x^{m-k}]\frac{m!d_m^{(r)}(x)}{2^m} - [x^{m-1-k}]\frac{(m-1)!d_{m-1}^{(r)}(x)}{2^{m-1}} \\ &= \frac{(m-1)!}{2^m}\Big([x^{m-k}]d_{m-1}^{(r)}(x) + (m-1+2r)[x^{m-k}]d_{m-2}^{(r)}(x)\Big). \end{aligned}$$

Hence,

$$\begin{aligned} &[x^{n-k}]\frac{n!d_n^{(r)}(x)}{2^n} - [x^0]\frac{k!d_k^{(r)}(x)}{2^k} \\ &= \sum_{m=k+1}^{n}\left([x^{m-k}]\frac{m!d_m^{(r)}(x)}{2^m} - [x^{m-1-k}]\frac{(m-1)!d_{m-1}^{(r)}(x)}{2^{m-1}}\right) \\ &= \sum_{m=k+1}^{n}\frac{(m-1)!}{2^m}\Big([x^{m-k}]d_{m-1}^{(r)}(x) + (m-1+2r)[x^{m-k}]d_{m-2}^{(r)}(x)\Big). \end{aligned}$$

Since $kd_k^{(r)}(x) = (1+2x)d_{k-1}^{(r)}(x) + (k-1+2r)d_{k-2}^{(r)}(x)$, we see that $[x^0]kd_k^{(r)}(x) = [x^0]d_{k-1}^{(r)}(x) + (k-1+2r)[x^0]d_{k-2}^{(r)}(x)$ and so

$$[x^{n-k}]\frac{n!d_n^{(r)}(x)}{2^n} = \sum_{m=k}^{n}\frac{(m-1)!}{2^m}\Big([x^{m-k}]d_{m-1}^{(r)}(x) + (m-1+2r)[x^{m-k}]d_{m-2}^{(r)}(x)\Big).$$

This yields the result.

Theorem 3.4. *Let $n \in \mathbb{Z}^+$. Then*

$$[x^n]d_n^{(r)}(x) = \frac{2^n}{n!},\ [x^{n-1}]d_n^{(r)}(x) = \frac{2^{n-1}}{(n-1)!},\quad [x^{n-2}]d_n^{(r)}(x) = \frac{2^{n-2}}{(n-2)!}\Big(r+\frac{n+1}{3}\Big)\ (n\ge 2),$$
$$[x^{n-3}]d_n^{(r)}(x) = \frac{2^{n-3}}{(n-3)!}\Big(r+\frac{n}{3}\Big)\ (n\ge 3),$$
$$[x^{n-4}]d_n^{(r)}(x) = \frac{2^{n-4}}{6\cdot(n-4)!}\Big(3r^2+(2n+1)r+\frac{(n+1)(5n-2)}{15}\Big)\ (n\ge 4).$$

Proof. Since $d_n(x) = \sum_{k=0}^n \binom{n}{k}\binom{x}{k}2^k$, we must have $[x^n]d_n(x) = \frac{2^n}{n!}$ and

$$[x^{n-1}]d_n(x) = [x^{n-1}]\Big(\binom{x}{n}2^n + n\binom{x}{n-1}2^{n-1}\Big) = \frac{2^n}{n!}\sum_{s=1}^{n-1}(-s) + \frac{2^{n-1}n}{(n-1)!} = \frac{2^{n-1}}{(n-1)!}.$$

By Theorem 3.3, $d_n^{(r)}(x) = \sum_{k=0}^n \binom{r-1+k}{k}d_{n-2k}(x)$. Hence

$$[x^n]d_n^{(r)}(x) = [x^n]d_n(x) = \frac{2^n}{n!}\quad\text{and}\quad [x^{n-1}]d_n^{(r)}(x) = [x^{n-1}]d_n(x) = \frac{2^{n-1}}{(n-1)!}.$$

This together with Lemma 3.1 (with $k=2$) yields for $n\ge 2$,

$$\begin{aligned}
[x^{n-2}]d_n^{(r)}(x) &= \sum_{m=2}^{n}\frac{2^{n-m}\cdot(m-1)!}{n!}\Big([x^{m-2}]d_{m-1}^{(r)}(x) + (m-1+2r)[x^{m-2}]d_{m-2}^{(r)}(x)\Big)\\
&= \sum_{m=2}^{n}\frac{2^{n-m}\cdot(m-1)!}{n!}\Big(\frac{2^{m-2}}{(m-2)!} + (m-1+2r)\frac{2^{m-2}}{(m-2)!}\\
&= \sum_{m=1}^{n}\frac{2^{n-2}}{n!}(m-1)(m+2r) = \frac{2^{n-2}}{n!}\sum_{m=1}^{n}(m^2+(2r-1)m-2r)\\
&= \frac{2^{n-2}}{n!}\Big(\frac{n(n+1)(2n+1)}{6} + (2r-1)\frac{n(n+1)}{2} - 2rn\Big)\\
&= \frac{2^{n-2}}{(n-2)!}\Big(r+\frac{n+1}{3}\Big).
\end{aligned}$$

Taking $k=3$ in Lemma 3.1 and then applying the above we deduce that for $n\ge 3$,

$$[x^{n-3}]d_n^{(r)}(x) = \sum_{m=3}^{n}\frac{2^{n-m}\cdot(m-1)!}{n!}\Big([x^{m-3}]d_{m-1}^{(r)}(x) + (m-1+2r)[x^{m-3}]d_{m-2}^{(r)}(x)\Big)$$

$$= \sum_{m=3}^{n} \frac{2^{n-m}\cdot(m-1)!}{n!}\Big(\frac{2^{m-3}}{(m-3)!}\Big(r+\frac{m}{3}\Big)+(m-1+2r)\frac{2^{m-3}}{(m-3)!}\Big)$$
$$= \sum_{m=3}^{n} \frac{2^{n-3}}{n!}(m-1)(m-2)\Big(3r-1+\frac{4m}{3}\Big)$$
$$= \frac{2^{n-3}}{n!}\Big\{8\sum_{m=3}^{n}\binom{m}{3}+(6r-2)\sum_{m=3}^{n}\binom{m-1}{2}\Big\}$$
$$= \frac{2^{n-3}}{n!}\Big\{8\binom{n+1}{4}+(6r-2)\binom{n}{3}\Big\} = \frac{2^{n-3}}{(n-3)!}\Big(r+\frac{n}{3}\Big),$$

where in the last step we use the identity $\sum_{m=r}^{n}\binom{m}{r}=\binom{n+1}{r+1}$. See [G,(1.48)]. Taking $k=4$ in Lemma 3.1 and then applying the above we deduce that for $n\geq 4$,

$$[x^{n-4}]d_n^{(r)}(x)$$
$$= \sum_{m=4}^{n} \frac{2^{n-m}\cdot(m-1)!}{n!}\Big([x^{m-4}]d_{m-1}^{(r)}(x)+(m-1+2r)[x^{m-4}]d_{m-2}^{(r)}(x)\Big)$$
$$= \sum_{m=4}^{n} \frac{2^{n-m}\cdot(m-1)!}{n!}\Big(\frac{2^{m-4}}{(m-4)!}\Big(r+\frac{m-1}{3}\Big)+(m-1+2r)\frac{2^{m-4}}{(m-4)!}\Big(r+\frac{m-1}{3}\Big)\Big)$$
$$= \sum_{m=4}^{n} \frac{2^{n-4}(m-1)(m-2)(m-3)}{n!}(m+2r)\Big(r+\frac{m-1}{3}\Big)$$
$$= \sum_{m=4}^{n} \frac{2^{n-4}(m-1)(m-2)(m-3)}{3\cdot n!}(m(m+1)+(5r-2)m+2r(3r-1))$$
$$= \frac{2^{n-4}}{n!}\Big\{40\sum_{m=4}^{n}\binom{m+1}{5}+8(5r-2)\sum_{m=4}^{n}\binom{m}{4}+4r(3r-1)\sum_{m=4}^{n}\binom{m-1}{3}\Big\}$$
$$= \frac{2^{n-4}}{n!}\Big\{40\binom{n+2}{6}+8(5r-2)\binom{n+1}{5}+4r(3r-1)\binom{n}{4}\Big\}$$
$$= \frac{2^{n-4}}{n!}\binom{n}{4}\Big\{40\cdot\frac{(n+2)(n+1)}{6\cdot 5}+8(5r-2)\frac{n+1}{5}+4r(3r-1)\Big\}$$
$$= \frac{2^{n-4}}{6\cdot(n-4)!}\Big(3r^2+(2n+1)r+\frac{(n+1)(5n-2)}{15}\Big).$$

This completes the proof.

Theorem 3.5. *For $n\in\mathbb{Z}^+$ we have*

(i) $d_n^{(r)}(x) = d_n^{(r+1)}(x) - d_{n-2}^{(r+1)}(x)$,

(ii) $d_n^{(r+1)}(x) = \sum_{k=0}^{[n/2]} d_{n-2k}^{(r)}(x)$,

(iii) $(n+1)^2 d_{n+1}^{(r)}(x)^2 - (n+2r+1)^2 d_n^{(r)}(x)^2 = 4(x-r)(x+1+r)(d_n^{(r+1)}(x)^2 - d_{n-1}^{(r+1)}(x)^2)$,

(iv) $(2r+1)\sum_{k=0}^{n-1}(2k+2r+1)d_k^{(r)}(x)^2 = n^2 d_n^{(r)}(x)^2 - 4(x-r)(x+1+r)d_{n-1}^{(r+1)}(x)^2$.

Proof. From Theorem 3.1 we know that for $|t|<1$,

$$(1-t^2)\sum_{m=0}^{\infty} d_m^{(r+1)}(x)t^m = \sum_{n=0}^{\infty} d_n^{(r)}(x)t^n$$

and

$$\sum_{n=0}^{\infty} d_n^{(r+1)}(x)t^n = \frac{1}{1-t^2}\sum_{m=0}^{\infty} d_m^{(r)}(x)t^m = \Big(\sum_{k=0}^{\infty} t^{2k}\Big)\Big(\sum_{m=0}^{\infty} d_m^{(r)}(x)t^m\Big).$$

Now comparing the coefficients of t^n on both sides we deduce (i) and (ii).

By (i),

$$\begin{aligned} d_{n+1}^{(r)}(x) &= d_{n+1}^{(r+1)}(x) - d_{n-1}^{(r+1)}(x) = \frac{(2x+1)d_n^{(r+1)}(x) + (n+2+2r)d_{n-1}^{(r+1)}(x)}{n+1} - d_{n-1}^{(r+1)}(x) \\ &= \frac{(2x+1)d_n^{(r+1)}(x) + (2r+1)d_{n-1}^{(r+1)}(x)}{n+1} \end{aligned}$$

and

$$\begin{aligned} d_n^{(r)}(x) &= d_n^{(r+1)}(x) - d_{n-2}^{(r+1)}(x) = d_n^{(r+1)}(x) - \frac{nd_n^{(r+1)}(x) - (2x+1)d_{n-1}^{(r+1)}(x)}{n+1+2r} \\ &= \frac{(2r+1)d_n^{(r+1)}(x) + (2x+1)d_{n-1}^{(r+1)}(x)}{n+1+2r}. \end{aligned}$$

Thus,

$$\begin{aligned} &(n+1)^2 d_{n+1}^{(r)}(x)^2 - (n+1+2r)^2 d_n^{(r)}(x)^2 \\ &= ((2x+1)d_n^{(r+1)}(x) + (2r+1)d_{n-1}^{(r+1)}(x))^2 - ((2r+1)d_n^{(r+1)}(x) + (2x+1)d_{n-1}^{(r+1)}(x))^2 \\ &= 4(x-r)(x+1+r)(d_n^{(r+1)}(x)^2 - d_{n-1}^{(r+1)}(x)^2). \end{aligned}$$

This proves (iii). By (iii),

$$\begin{aligned} &\sum_{k=0}^{n-1}(2r+1)(2k+2r+1)d_k^{(r)}(x)^2 \\ &= \sum_{k=0}^{n-1}\big((k+1)^2 d_{k+1}^{(r)}(x)^2 - k^2 d_k^{(r)}(x)^2\big) - 4(x-r)(x+1+r)\sum_{k=0}^{n-1}\big(d_k^{(r+1)}(x)^2 - d_{k-1}^{(r+1)}(x)^2\big) \\ &= n^2 d_n^{(r)}(x)^2 - 4(x-r)(x+1+r)d_{n-1}^{(r+1)}(x)^2. \end{aligned}$$

This proves (iv). Thus the theorem is proved.

Theorem 3.6. *Let $n\in\mathbb{Z}^+$, $r\in\{0,1,2,\ldots\}$ and $x\in\mathbb{Z}$. Then*

$$\begin{aligned}&(2r+1)\prod_{k=-r}^{r}(x+k)(x+1-k)\cdot\sum_{k=0}^{n-1}(2k+2r+1)d_k^{(r)}(x)^2\\&\equiv 0\ (\mathrm{mod}\ 2n^2(n+1)^2\cdots(n+2r)^2).\end{aligned}$$

Proof. By Theorem 3.5(iv) and (3.6),

$$\begin{aligned}&(2r+1)\prod_{k=-r}^{r}(x+k)(x+1-k)\sum_{k=0}^{n-1}(2k+2r+1)d_k^{(r)}(x)^2\\&=\prod_{k=-r}^{r}(x+k)(x+1-k)\cdot\big(n^2d_n^{(r)}(x)^2-4(x-r)(x+1+r)d_{n-1}^{(r+1)}(x)^2\big)\\&=(x-r)(x+r+1)(n+2r)^2(n+2r-1)^2\cdots(n+1)^2n^2\Big(\sum_{k=0}^{n}\binom{n}{k}\binom{x+r+k}{n+2r}\Big)^2\\&\quad-4(n+2r+1)^2(n+2r)^2\cdots n^2\Big(\sum_{k=0}^{n-1}\binom{n-1}{k}\binom{x+r+1+k}{n+2r+1}\Big)^2.\end{aligned}$$

Note that $(x+r+1)(x-r)\equiv 0\ (\mathrm{mod}\ 2)$. From the above we deduce the result.

We remark that Theorem 3.6 is a vast generalization of (1.2), and the next theorem is a vast generalization of (1.4).

Theorem 3.7. *Suppose $r\notin\{-\frac12,-\frac22,-\frac32,\ldots\}$. Then*

$$d_n^{(r)}(x)^2=\binom{n+2r}{n}\sum_{m=0}^{n}\frac{\binom{x-r}{m}\binom{x+r+m}{m}\binom{n+2r+m}{n-m}}{\binom{m+2r}{m}}4^m\quad\textit{for}\quad n=0,1,2,\ldots.$$

For $r\in\{0,1,2,\ldots\}$ we also have

$$\binom{x+r}{2r}d_n^{(r)}(x)^2=\binom{n+2r}{2r}\sum_{m=0}^{n}\binom{n+2r}{m+2r}\binom{n+2r+m}{m}\binom{x+m+r}{2m+2r}4^m\ (n=0,1,2,\ldots).$$

Proof. Set

$$s(n)=\frac{d_n^{(r)}(x)^2}{\binom{n+2r}{n}}\quad\text{and}\quad S(n)=\sum_{m=0}^{n}\frac{\binom{x-r}{m}\binom{x+r+m}{m}\binom{n+2r+m}{n-m}}{\binom{m+2r}{m}}4^m.$$

Using Maple we find that for $n=1,2,3,\ldots$,

$$(n+2)(n+2+2r)S(n+2)-((2x+1)^2+(n+1)(n+1+2r))(S(n+1)+S(n))+n(n+2r)S(n-1)=0.$$

By Theorem 3.2,

$$d_{n+2}^{(r)}(x)=\frac{(1+2x)d_{n+1}^{(r)}(x)+(n+1+2r)d_n^{(r)}(x)}{n+2},\ d_{n-1}^{(r)}(x)=\frac{(n+1)d_{n+1}^{(r)}(x)-(1+2x)d_n^{(r)}(x)}{n+2r}.$$

Thus,

$$\begin{aligned}
&(n+2)(n+2+2r)s(n+2)+n(n+2r)s(n-1)\\
&=\frac{(n+2)(n+2+2r)}{\binom{n+2+2r}{2r}}d_{n+2}^{(r)}(x)^2+\frac{n(n+2r)}{\binom{n-1+2r}{2r}}d_{n-1}^{(r)}(x)^2\\
&=\frac{((1+2x)d_{n+1}^{(r)}(x)+(n+1+2r)d_n^{(r)}(x))^2}{\binom{n+1+2r}{2r}}+\frac{((n+1)d_{n+1}^{(r)}(x)-(1+2x)d_n^{(r)}(x))^2}{\binom{n+2r}{2r}}\\
&=d_{n+1}^{(r)}(x)^2\Big\{\frac{(1+2x)^2}{\binom{n+1+2r}{2r}}+\frac{(n+1)^2}{\binom{n+2r}{2r}}\Big\}+d_n^{(r)}(x)^2\Big\{\frac{(1+2x)^2}{\binom{n+2r}{2r}}+\frac{(n+1+2r)^2}{\binom{n+1+2r}{2r}}\Big\}\\
&=\frac{d_{n+1}^{(r)}(x)^2}{\binom{n+1+2r}{2r}}\big((1+2x)^2+(n+1)(n+1+2r)\big)+\frac{d_n^{(r)}(x)^2}{\binom{n+2r}{2r}}\big((1+2x)^2+(n+1)(n+1+2r)\big).\\
&=((1+2x)^2+(n+1)(n+1+2r))(s(n)+s(n+1)).
\end{aligned}$$

This shows that $s(n)$ and $S(n)$ satisfy the same recurrence relation. Also,

$$s(0)=1=S(0),\ s(1)=\frac{(1+2x)^2}{2r+1}=S(1),\ s(2)=\frac{(2x^2+2x+r+1)^2}{(r+1)(2r+1)}=S(2).$$

Thus, $s(n)=S(n)$ for $n=0,1,2,\ldots$. To finish the proof, we note that for $r=0,1,2,\ldots$,

$$\begin{aligned}
&\binom{x+r}{2r}\binom{x-r}{m}\binom{x+r+m}{m}\binom{n+2r+m}{2m+2r}\\
&=\binom{m+2r}{2r}\binom{n+2r}{m+2r}\binom{n+2r+m}{m}\binom{x+m+r}{2m+2r}.
\end{aligned}$$

Many orthogonal polynomials have formulas for linearization of their products. As examples, for Hermite polynomials $\{H_n(x)\}$ ($H_{-1}(x)=0,\ H_0(x)=1,\ H_{n+1}(x)=2xH_n(x)-2nH_{n-1}(x)$) we have

$$H_m(x)H_n(x)=\sum_{k=0}^{\min\{m,n\}}\binom{m}{k}\binom{n}{k}2^k k!H_{m+n-2k}(x);$$

for Legendre polynomials $\{P_n(x)\}$ we have

$$P_m(x)P_n(x)=\sum_{k=0}^{\min\{m,n\}}\frac{2m+2n+1-4k}{2m+2n+1-2k}\cdot\frac{\binom{2k}{k}\binom{2m-2k}{m-k}\binom{2n-2k}{n-k}}{\binom{2(m+n-k)}{m+n-k}}P_{m+n-2k}(x).$$

See [AAR, Theorem 6.8.1 and Corollary 6.8.3].

The author is lucky to discover the following linearization of $d_m^{(r)}(x)d_n^{(r)}(x)$:

Theorem 3.8. *Let m and n be nonnegative integers. Then*

$$d_m^{(r)}(x)d_n^{(r)}(x)=\sum_{k=0}^{\min\{m,n\}}\binom{m+n-2k}{m-k}\binom{2r+m+n-k}{k}(-1)^k d_{m+n-2k}^{(r)}(x).$$

By the definition of $d_n^{(r)}(x)$, Theorem 3.8 is equivalent to the following identity:

$$\sum_{k=0}^{m}\sum_{l=0}^{n}\binom{x+r+k}{k}\binom{x+r+l}{l}\binom{x-r}{m-k}\binom{x-r}{n-l}$$
$$=\sum_{k=0}^{m}\sum_{l=0}^{m+n-2k}\binom{m+n-2k}{m-k}\binom{2r+m+n-k}{k}(-1)^k\binom{x+r+l}{l}\binom{x-r}{m+n-2k-l}.$$

By the Maple package DoubleSum (see http://cam.tju.edu.cn/∼hou/soft/ds.html and the method in [CHM]) one may prove the above identity.

Conjecture 3.1. *Let $n\in\mathbb{Z}^+$, $r\ge 0$ and $x\in(-1,0)$. Then*

$$(-1)^n(d_n^{(r)}(x)^2-d_{n+1}^{(r)}(x)d_{n-1}^{(r)}(x))>0.$$

4. The orthogonal polynomials $\{D_n^{(r)}(x)\}$

In this section we discuss a kind of orthogonal polynomials related to $\{d_n^{(r)}(x)\}$.

Definition 4.1. *Let $\{D_n^{(r)}(x)\}$ be the polynomials given by*

$$D_0^{(r)}(x)=1,\ D_1^{(r)}(x)=x\quad\text{and}\quad D_{n+1}^{(r)}(x)=xD_n^{(r)}(x)-n(n+2r)D_{n-1}^{(r)}(x)\ (n\ge 1).$$

By [GR, p.175] or [AAR, pp.244-245], for $r>-\frac{1}{2}$, $\{D_n^{(r)}(x)\}$ are orthogonal polynomials. More precisely, for given $r>-\frac{1}{2}$ there is a weight function $w(x)$ such that

$$\int_{-\infty}^{\infty}w(x)D_m^{(r)}(x)D_n^{(r)}(x)dx=\begin{cases}(2r+1)^2(2r+2)^2\cdots(2r+n)^2 & \text{if } m=n,\\ 0 & \text{if } m\ne n.\end{cases}$$

The first few $D_n^{(r)}(x)$ are shown below:

$$D_0^{(r)}(x)=1,\ D_1^{(r)}(x)=x,\ D_2^{(r)}(x)=x^2-2r-1,\ D_3^{(r)}(x)=x^3-(6r+5)x,$$
$$D_4^{(r)}(x)=x^4-(12r+14)x^2+12r^2+24r+9,$$
$$D_5^{(r)}(x)=x^5-(20r+30)x^3+(60r^2+160r+89)x.$$

Lemma 4.1. *For any nonnegative n we have*

$$d_n^{(r)}(x)=\frac{i^nD_n^{(r)}(-i(1+2x))}{n!}\quad\text{and so}\quad D_n^{(r)}(x)=(-i)^nn!d_n^{(r)}\Big(\frac{ix-1}{2}\Big).$$

Proof. Set $p_n^{(r)}(x)=\frac{i^nD_n^{(r)}(-i(1+2x))}{n!}$. Then $p_0^{(r)}(x)=1=d_0^{(r)}(x)$ and $p_1^{(r)}(x)=1+2x=d_1^{(r)}(x)$. As

$$(n+1)p_n^{(r)}(x)=\frac{i^{n+1}D_{n+1}^{(r)}(-i(1+2x))}{n!}$$

$$= \frac{i^{n+1}}{n!}\big(- i(1+2x)D_n^{(r)}(-i(1+2x)) - n(n+2r)D_{n-1}^{(r)}(-i(1+2x))\big)$$
$$= (1+2x)p_n^{(r)}(x) + (n+2r)p_{n-1}^{(r)}(x),$$

we must have $p_n^{(r)}(x) = d_n^{(r)}(x)$, which yields the result.

Let $\Gamma(x)$ be the Gamma function defined by $\Gamma(x) = \int_0^{+\infty} t^{x-1}\mathrm{e}^{-t}dt$ for $x > 0$. It is well known that $\Gamma(x+1) = x\Gamma(x)$ and so $\Gamma(n) = (n-1)!$ for $n \in \mathbb{Z}^+$.

Theorem 4.1. *For any positive integer n and $r > -\frac{1}{2}$ we have*

$$\sum_{k=0}^{n-1}(2k+2r+1)\frac{D_k^{(r)}(x)^2}{k!\Gamma(k+2r+1)} = \frac{D_n^{(r)}(x)^2 - D_{n-1}^{(r)}(x)D_{n+1}^{(r)}(x)}{(n-1)!\Gamma(n+2r)}$$

and so

$$D_n^{(r)}(x)^2 - D_{n+1}^{(r)}(x)D_{n-1}^{(r)}(x) > 0 \quad \textit{for real} \quad x.$$

Proof. It is well known that $\Gamma(t+1) = t\Gamma(t)$ and $\Gamma(t) > 0$ for $t > 0$. Thus,

$$\begin{aligned}
&\frac{D_{n+1}^{(r)}(x)^2 - D_n^{(r)}(x)D_{n+2}^{(r)}(x)}{n!\Gamma(n+2r+1)} - \frac{D_n^{(r)}(x)^2 - D_{n-1}^{(r)}(x)D_{n+1}^{(r)}(x)}{(n-1)!\Gamma(n+2r)}\\
&= \frac{1}{n!\Gamma(n+2r+1)}\Big(D_{n+1}^{(r)}(x)^2 - D_n^{(r)}(x)(xD_{n+1}^{(r)}(x) - (n+1)(n+2r+1)D_n^{(r)}(x))\\
&\quad - n(n+2r)(D_n^{(r)}(x)^2 - D_{n-1}^{(r)}(x)D_{n+1}^{(r)}(x))\Big)\\
&= \frac{1}{n!\Gamma(n+2r+1)}\big(D_{n+1}^{(r)}(x)(D_{n+1}^{(r)}(x) - xD_n^{(r)}(x) + n(n+2r)D_{n-1}^{(r)}(x))\\
&\quad + ((n+1)(n+1+2r) - n(n+2r))D_n^{(r)}(x)^2\big)\\
&= (2n+2r+1)\frac{D_n^{(r)}(x)^2}{n!\Gamma(n+2r+1)}.
\end{aligned}$$

By the above, the identity can be easily proved by induction on n.

Corollary 4.1. *Let $n \in \mathbb{Z}^+$ and $r > -\frac{1}{2}$. Then*

$$\sum_{k=0}^{n-1}(-1)^k(2k+2r+1)\frac{k!d_k^{(r)}(x)^2}{\Gamma(k+2r+1)} = \frac{(-1)^n n!}{\Gamma(n+2r)}\big(nd_n^{(r)}(x)^2 - (n+1)d_{n-1}^{(r)}(x)d_{n+1}^{(r)}(x)\big).$$

In particular, for $r = 0$ we have

$$\sum_{k=0}^{n-1}(-1)^k(2k+1)d_k(x)^2 = (-1)^n n(nd_n(x)^2 - (n+1)d_{n-1}(x)d_{n+1}(x)).$$

Proof. Replacing x with $-i(1+2x)$ in Theorem 4.1 and then applying Lemma 4.1 yields the result.

Theorem 4.2. *Let n be a positive integer and $r > -\frac{1}{2}$. Then*

$$\sum_{k=0}^{n-1}\frac{D_k^{(r)}(x)^2}{k!\Gamma(k+2r+1)} = \frac{D_{n-1}^{(r)}(x)\frac{d}{dx}D_n^{(r)}(x) - D_n^{(r)}(x)\frac{d}{dx}D_{n-1}^{(r)}(x)}{(n-1)!\Gamma(n+2r)}$$

and

$$\sum_{k=0}^{n-1}\frac{(-1)^k k!}{\Gamma(k+2r+1)}d_k^{(r)}(x)^2=\frac{(-1)^{n-1}n!}{2\Gamma(n+2r)}\big(d_{n-1}^{(r)}(x)\frac{d}{dx}d_n^{(r)}(x)-d_n^{(r)}(x)\frac{d}{dx}d_{n-1}^{(r)}(x)\big).$$

Proof. Since $D_{n+1}^{(r)}(x)=xD_n^{(r)}(x)-n(n+2r)D_{n-1}^{(r)}(x)$ we see that

$$\frac{d}{dx}D_{n+1}^{(r)}(x)=D_n^{(r)}(x)+x\frac{d}{dx}D_n^{(r)}(x)-n(n+2r)\frac{d}{dx}D_{n-1}^{(r)}(x).$$

As $\Gamma(t+1)=t\Gamma(t)$, we deduce that

$$\begin{aligned}
&\frac{D_n^{(r)}(x)\frac{d}{dx}D_{n+1}^{(r)}(x)-D_{n+1}^{(r)}(x)\frac{d}{dx}D_n^{(r)}(x)}{n!\Gamma(n+2r+1)}-\frac{D_{n-1}^{(r)}(x)\frac{d}{dx}D_n^{(r)}(x)-D_n^{(r)}(x)\frac{d}{dx}D_{n-1}^{(r)}(x)}{(n-1)!\Gamma(n+2r)}\\
&=\frac{1}{n!\Gamma(n+2r+1)}\Big\{D_n^{(r)}(x)^2+xD_n^{(r)}(x)\frac{d}{dx}D_n^{(r)}(x)-n(n+2r)D_n^{(r)}(x)\frac{d}{dx}D_{n-1}^{(r)}(x)\\
&\quad-(xD_n^{(r)}(x)-n(n+2r)D_{n-1}^{(r)}(x))\frac{d}{dx}D_n^{(r)}(x)\\
&\quad-n(n+2r)D_{n-1}^{(r)}(x)\frac{d}{dx}D_n^{(r)}(x)+n(n+2r)D_n^{(r)}(x)\frac{d}{dx}D_{n-1}^{(r)}(x)\Big\}\\
&=\frac{D_n^{(r)}(x)^2}{n!\Gamma(n+2r+1)}.
\end{aligned}$$

Thus the first identity is easily proved by induction on n.

By Lemma 4.1, $d_n^{(r)}(x)=i^nD_n^{(r)}(-i(1+2x))/n!$. Thus, $\frac{d}{dx}d_n^{(r)}(x)=i^n\frac{d}{dx}D_n^{(r)}(-i(1+2x))(-2i)/n!$. Hence applying the above we obtain

$$\begin{aligned}
&\sum_{k=0}^{n-1}\frac{(-1)^k k!}{\Gamma(k+2r+1)}d_k^{(r)}(x)^2\\
&=\sum_{k=0}^{n-1}\frac{D_k^{(r)}(-i(1+2x))^2}{k!\Gamma(k+2r+1)}\\
&=\frac{\frac{d}{dx}D_n^{(r)}(-i(1+2x))\cdot D_{n-1}^{(r)}(-i(1+2x))-D_n^{(r)}(-i(1+2x))\frac{d}{dx}D_{n-1}^{(r)}(-i(1+2x))}{(n-1)!\Gamma(n+2r)}\\
&=\frac{1}{(n-1)!\Gamma(n+2r)}\Big(\frac{n!\frac{d}{dx}d_n^{(r)}(x)}{(-2i)i^n}\cdot\frac{(n-1)!d_{n-1}^{(r)}(x)}{i^{n-1}}-\frac{n!d_n^{(r)}(x)}{i^n}\cdot\frac{(n-1)!\frac{d}{dx}d_{n-1}^{(r)}(x)}{(-2i)i^{n-1}}\Big)\\
&=\frac{(-1)^{n-1}n!}{2\Gamma(n+2r)}\big(d_{n-1}^{(r)}(x)\frac{d}{dx}d_n^{(r)}(x)-d_n^{(r)}(x)\frac{d}{dx}d_{n-1}^{(r)}(x)\big).
\end{aligned}$$

This completes the proof.

Theorem 4.3. *For any nonnegative integer n we have*

$$D_n^{(r)}(x)^2=\sum_{m=0}^{n}\binom{n+2r+m}{n-m}(-1)^{n-m}\prod_{j=m+1}^{n}j(2r+j)\prod_{k=1}^{m}(x^2+(2r+2k-1)^2).$$

Proof. By Lemma 4.1 and Theorem 3.7,

$$D_n^{(r)}(x)^2=(-1)^n n!^2 d_n^{(r)}\Big(\frac{ix-1}{2}\Big)^2=(-1)^n n!^2\binom{n+2r}{n}\sum_{m=0}^{n}\frac{\binom{\frac{ix-1}{2}-r}{m}\binom{\frac{ix-1}{2}+r+m}{m}\binom{n+2r+m}{n-m}}{\binom{m+2r}{m}}4^m.$$

Since

$$\begin{aligned}&\binom{\frac{ix-1}{2}-r}{m}\binom{\frac{ix-1}{2}+r+m}{m}\\&=\frac{(\frac{ix-1}{2}-r)(\frac{ix-1}{2}-(r+1))\cdots(\frac{ix-1}{2}-(r+m-1))(\frac{ix-1}{2}+r+m)\cdots(\frac{ix-1}{2}+r+1)}{m!^2}\\&=\frac{((ix)^2-(2r+1)^2)\cdots((ix)^2-(2r+2m-1)^2)}{2^{2m}\cdot m!^2}=\frac{(x^2+(2r+1)^2)\cdots(x^2+(2r+2m-1)^2)}{(-4)^m\cdot m!^2},\end{aligned}$$

from the above we deduce that

$$\begin{aligned}D_n^{(r)}(x)^2&=(-1)^n n!(2r+1)(2r+2)\cdots(2r+n)\\&\quad\times\sum_{m=0}^{n}\binom{n+2r+m}{n-m}\frac{(-1)^m}{m!(2r+1)(2r+2)\cdots(2r+m)}\prod_{k=1}^{m}(x^2+(2r+2k-1)^2).\end{aligned}$$

This yields the result.

Theorem 4.4. *We have*

$$\sum_{n=0}^{\infty}D_n^{(r)}(x)\frac{t^n}{n!}=(1+t^2)^{-r-\frac{1}{2}}\ \mathrm{e}^{x\ arctan\ t}$$

and so

$$\sum_{n=0}^{\infty}D_n^{(r)}(x)\frac{\tan^n\theta}{n!}=\mathrm{e}^{x\theta}(\cos\ \theta)^{2r+1}\ (0<\theta<\frac{\pi}{2}).$$

Proof. Set $f(t)=\sum_{n=0}^{\infty}\ D_n^{(r)}(x)\frac{t^n}{n!}$. Then

$$f(t)=1+\sum_{n=0}^{\infty}D_{n+1}^{(r)}(x)\frac{t^{n+1}}{(n+1)!}=1+\sum_{n=0}^{\infty}xD_n^{(r)}(x)\frac{t^{n+1}}{(n+1)!}-\sum_{n=1}^{\infty}n(n+2r)D_{n-1}^{(r)}(x)\frac{t^{n+1}}{(n+1)!}.$$

Hence

$$\begin{aligned}f'(t)&=\sum_{n=0}^{\infty}xD_n^{(r)}(x)\frac{t^n}{n!}-\sum_{n=1}^{\infty}(n+2r)D_{n-1}^{(r)}(x)\frac{t^n}{(n-1)!}\\&=xf(t)-2rtf(t)-t\Big(\sum_{n=1}^{\infty}D_{n-1}^{(r)}(x)\frac{t^n}{(n-1)!}\Big)'\\&=(x-2rt)f(t)-t(tf(t))'=(x-2rt)f(t)-t(f(t)+tf'(t)).\end{aligned}$$

That is,

$$\frac{f'(t)}{f(t)}=\frac{x-(2r+1)t}{1+t^2}.$$

Thus,

$$\int \frac{f'(t)}{f(t)}dt = x\int \frac{dt}{1+t^2} - \frac{2r+1}{2}\int \frac{d(1+t^2)}{1+t^2}\Big) = x\ \arctan\ t - \frac{2r+1}{2}\log(1+t^2) + C_1.$$

Hence

$$\log f(t) = x\ \arctan\ t - \frac{2r+1}{2}\log(1+t^2) + C_2 \quad \text{and so} \quad f(t) = C(1+t^2)^{-r-\frac{1}{2}}\ \mathrm{e}^{x\ arctan\ t}.$$

As $f(0) = 1$ we see that $C = 1$. So the theorem is proved.

Corollary 4.2. *For any nonnegative integer n,*

$$D_n^{(r)}(-x) = (-1)^n D_n^{(r)}(x) \quad \textit{and} \quad D_n^{(r)}(0) = \begin{cases} 0 & \textit{if } n \textit{ is odd,} \\ n!\binom{-r-\frac{1}{2}}{\frac{n}{2}} & \textit{if } n \textit{ is even.} \end{cases}$$

Proof. By Theorem 4.4,

$$\sum_{n=0}^{\infty} D_n^{(r)}(-x)\frac{(-t)^n}{n!} = (1+t^2)^{-r-\frac{1}{2}}\ \mathrm{e}^{-x\ arctan\ (-t)} = (1+t^2)^{-r-\frac{1}{2}}\ \mathrm{e}^{x\ arctan\ t} = \sum_{n=0}^{\infty} D_n^{(r)}(x)\frac{t^n}{n!}.$$

Thus, $(-1)^n D_n^{(r)}(-x) = D_n^{(r)}(x)$. Taking $x = 0$ in Theorem 4.4 and then applying Newton's binomial theorem we see that

$$\sum_{n=0}^{\infty} D_n^{(r)}(0)\frac{t^n}{n!} = (1+t^2)^{-r-\frac{1}{2}} = \sum_{k=0}^{\infty}\binom{-r-\frac{1}{2}}{k}t^{2k}.$$

Comparing the coefficients of t^n on both sides yields the remaining part.

Theorem 4.5. *For $n = 0, 1, 2, \ldots$ we have*

$$D_n^{(r)}(x) = x^n - \sum_{k=1}^{n-1} k(k+2r)D_{k-1}^{(r)}(x)x^{n-1-k}$$

and

$$n!d_n^{(r)}(x) = (1+2x)^n + \sum_{k=1}^{n-1}(k+2r)\cdot k!d_{k-1}^{(r)}(x)(1+2x)^{n-1-k}.$$

Proof. For $k = 0, 1, 2, \ldots$ we have

$$\frac{D_{k+1}^{(r)}(x)}{x^{k+1}} - \frac{D_k^{(r)}(x)}{x^k} = -k(k+2r)\frac{D_{k-1}^{(r)}(x)}{x^{k+1}}.$$

Thus,

$$-\sum_{k=1}^{n-1} k(k+2r)\frac{D_{k-1}^{(r)}(x)}{x^{k+1}} = \sum_{k=1}^{n-1}\Big(\frac{D_{k+1}^{(r)}(x)}{x^{k+1}} - \frac{D_k^{(r)}(x)}{x^k}\Big) = \frac{D_n^{(r)}(x)}{x^n} - \frac{D_1^{(r)}(x)}{x}.$$

Multiplying by x^n on both sides and noting that $D_1^{(r)}(x) = x$ we deduce the formula for $D_n^{(r)}(x)$.

By Lemma 4.1, $(-i)^n n! d_n^{(r)}(x) = D_n^{(r)}(-i(1+2x))$. Thus,

$$\begin{aligned}
&(-i)^n n! d_n^{(r)}(x) \\
&= D_n^{(r)}(-i(1+2x)) = (-i(1+2x))^n - \sum_{k=1}^{n-1} k(k+2r) D_{k-1}^{(r)}(-i(1+2x))(-i(1+2x))^{n-1-k} \\
&= (-i(1+2x))^n - \sum_{k=1}^{n-1} k(k+2r)(-i)^{k-1}(k-1)! d_{k-1}^{(r)}(x)(-i(1+2x))^{n-1-k} \\
&= (-i)^n \Big\{ (1+2x)^n + \sum_{k=1}^{n-1} (k+2r) \cdot k! d_{k-1}^{(r)}(x)(1+2x)^{n-1-k} \Big\}.
\end{aligned}$$

This completes the proof.

Corollary 4.3. *Let n be a positive integer. Then*

$$[x^n] D_n^{(r)}(x) = 1 \quad \textit{and} \quad [x^{n-2}] D_n^{(r)}(x) = -\frac{(n-1)n(2n-1+6r)}{6} \ (n \geq 2),$$

where $[x^k] f(x)$ is the coefficient of x^k in the power series expansion of $f(x)$.

Proof. From Theorem 4.5 we see that $[x^n] D_n^{(r)}(x) = 1$ and so

$$[x^{n-2}] D_n^{(r)}(x) = -\sum_{k=1}^{n-1} k(k+2r) = -\sum_{k=1}^{n-1} k^2 - 2r \sum_{k=1}^{n-1} k = -\frac{(n-1)n(2n-1)}{6} - rn(n-1).$$

This yields the result.

Theorem 4.6. *For $n = 2, 3, 4, \ldots$ we have*

$$D_n^{(r)}(x) = D_n^{(r+1)}(x) + n(n-1) D_{n-2}^{(r+1)}(x).$$

Proof. By Theorem 4.4,

$$(1+t^2) \sum_{n=0}^{\infty} D_n^{(r+1)}(x) \frac{t^n}{n!} = \sum_{n=0}^{\infty} D_n^{(r)}(x) \frac{t^n}{n!}.$$

Now comparing the coefficients of t^n on both sides we obtain $\frac{D_n^{(r)}(x)}{n!} = \frac{D_n^{(r+1)}(x)}{n!} + \frac{D_{n-2}^{(r+1)}(x)}{(n-2)!}$. This yields the result.

Theorem 4.7. *Let m and n be nonnegative integers. Then*

$$D_m^{(r)}(x) D_n^{(r)}(x) = \sum_{k=0}^{\min\{m,n\}} \binom{2r+m+n-k}{k} \frac{m!n!}{(m-k)!(n-k)!} D_{m+n-2k}^{(r)}(x).$$

Proof. This is immediate from Theorem 3.8 and Lemma 4.1.

Remark 4.1 Since $\sum_{k=0}^{n} \binom{n}{k}(-1)^k \binom{x}{k}(-2)^k = d_n(x)$, applying the binomial inversion formula we get $\sum_{k=0}^{n} \binom{n}{k}(-1)^k d_k(x) = \binom{x}{n}(-2)^n$. Recall that $D_k^{(0)}(x) = k!(-i)^k d_k\big(\frac{ix-1}{2}\big)$. From the above we deduce that

$$\sum_{k=0}^{n} \binom{n}{k} \frac{(-i)^k}{k!} D_k^{(0)}(x) = \binom{\frac{ix-1}{2}}{n} (-2)^n.$$

5. The orthogonal polynomials $\{C_n(x)\}$

Definition 5.1. *For any nonnegative integer n define*

$$C_n(x)=\sum_{k=0}^n\binom{n}{k}\frac{1}{k!^2}\prod_{j=0}^{k-1}(x-j(j+1))\quad\text{and}\quad c_n(x)=\sum_{k=0}^n\binom{n}{k}\binom{x}{k}\binom{x+k}{k}.$$

We note that $\{c_n(x)\}$ have been studied by Z.W. Sun[Su] and Guo[Gu]. Since

$$\begin{aligned}c_n(x)&=\sum_{k=0}^n\binom{n}{k}\binom{2k}{k}\binom{x+k}{2k}=\sum_{k=0}^n\binom{n}{k}\binom{2k}{k}\frac{1}{(2k)!}\prod_{j=0}^{k-1}(x-j)(x+j+1)\\&=\sum_{k=0}^n\binom{n}{k}\frac{1}{k!^2}\prod_{j=0}^{k-1}(x(x+1)-j(j+1)),\end{aligned}$$

we have

$$c_n(x)=C_n(x(x+1))\quad\text{and so}\quad C_n(x)=c_n\Big(\frac{-1+\sqrt{1+4x}}{2}\Big).\tag{5.1}$$

Theorem 5.1. *For $n\in\mathbb{Z}^+$ we have*

$$(n+1)^2C_{n+1}(x)=(x+1+2n(n+1))C_n(x)-n^2C_{n-1}(x).$$

Proof. Using Maple software and Zeilberger's algorithm we find that

$$(n+1)^2c_{n+1}(x)=(1+x(x+1)+2n(n+1))c_n(x)-n^2c_{n-1}(x).\tag{5.2}$$

Now replacing x with $\frac{-1+\sqrt{1+4x}}{2}$ and then applying (5.1) we deduce the result.

By [GR, p.175] and Theorem 5.1, $\{C_n(x)\}$ are orthogonal polynomials. The first few $\{C_n(x)\}$ are shown below:

$$C_0(x)=1,\ C_1(x)=x+1,\ C_2(x)=\frac{1}{4}x^2+\frac{3}{2}x+1,\ C_3(x)=\frac{1}{36}x^3+\frac{19}{36}x^2+\frac{11}{6}x+1,$$
$$C_4(x)=\frac{1}{576}x^4+\frac{11}{144}x^3+\frac{115}{144}x^2+\frac{25}{12}x+1,$$
$$C_5(x)=\frac{1}{14400}x^5+\frac{17}{2880}x^4+\frac{251}{1800}x^3+\frac{3799}{3600}x^2+\frac{137}{60}x+1.$$

Corollary 5.1. *For $n\in\mathbb{Z}^+$ we have*

$$\sum_{m=0}^{n-1}C_m(x)C_m(y)=\frac{n^2(C_n(x)C_{n-1}(y)-C_{n-1}(x)C_n(y))}{x-y}$$

and

$$\sum_{m=0}^{n-1}(2+x+y+4m(m+1))(-1)^{n-1-m}C_m(x)C_m(y)$$

$$= n^2(C_n(x)C_{n-1}(y) + C_{n-1}(x)C_n(y)).$$

Proof. Taking $a_n = n^2$, $b_n = 1$, $b'_n = 2n(n+1)$, $c = -1$ and $s_n(x) = C_n(x)$ in Theorem 2.2 and then applying Theorem 5.1 yields the result.

Let $\{z_n\}_{n\geq 0}$ be a sequence of positive numbers. It is called log-convex if $z_n^2 - z_{n-1}z_{n+1} \leq 0$ for all $n \in \mathbb{Z}^+$ and strictly log-convex if the inequality is strict. The sequence is called log-concave if the inequality changes its direction. Many known sequences have been proved to be log-convex or log-concave. As examples, the Apéry numbers $\{A_n\}$ is log-convex, and the Legendre polynomials $\{P_n(x)\}$ is log-concave when $|x| < 1$. See [LW], [XY] and [AAR, p.342].

Conjecture 5.1. *Let $n \in \mathbb{Z}^+$. If $x > 0$ or $x \leq -1$, then $C_n(x)^2 - C_{n-1}(x)C_{n+1}(x) > 0$. If $-\frac{1}{3} < x < 0$, then $C_n(x)^2 - C_{n-1}(x)C_{n+1}(x) < 0$.*

References

[AAR] G. Andrews, R. Askey and R. Roy, *Special Functions*, Encyclopedia of Mathematical Applications, Vol.71, Cambridge University Press, Cambridge, 1999.

[BS] C. Banderier and S. Schwer, *Why Delannoy numbers?*, J. Stat. Planning Inference **135**(2005), 40-54.

[Be] F. Beukers,*Another congruence for the Apéry numbers*, J. Number Theory **25**(1987), 201-210.

[CHM] W.Y.C. Chen, Q.-H. Hou and Y.-P. Mu, *A telescoping method for double summations*, J. Comput. Appl. Math. **196**(2006), 553-566.

[EMOT] A. Erdélyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, *Higher Transcendental Functions, Vol.II*, McGraw-Hill, New York, 1953. Based, in part, on notes left by Harry Bateman.

[GR] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, 2nd edition, Encyclopedia of Mathematics and its Applications, Vol.96, Cambridge University Press, Cambridge, 2004.

[G] H.W. Gould, Combinatorial Identities, A Standardized Set of Tables Listing 500 Binomial Coefficient Summations, West Virginia University, Morgantown, WV, 1972.

[Gu] V.J.W. Guo, *Proof of Sun's conjectures on integer-valued polynomials*, J. Math. Anal. Appl., 2016, in press, http://dx.doi.org/10.1016/j.jmaa.2016.06.028.

[LW] L.L. Liu and Y. Wang, *On the log-convexity of combinatorial sequences*, Adv. Appl. Math. **39**(2007), 453-476.

[Sl] L.J. Slater, *Generalized Hypergeometric Functions,* Cambridge University Press, Cambridge, 1966.

[Su] Z.W. Sun, *Supercongruences involving dual sequences*, preprint, arXiv:1512.00712v3, 2015.

[XY] E.X.W. Xia and O.X.M. Yao, *A criterion for the log-convexity of combinatorial sequences*, Electron. J. Combin. **20**(2013), #P3, 1-10.